

GLOBAL JOURNAL OF ENGINEERING SCIENCE AND RESEARCHES

RICCI RECURRENT PROJECTIVE TRANSFORMATION

B. K. Singh

Department of Mathematics, Amity University, Chhattisgarh

I. INTRODUCTION

Let F_n be an n-dimensional Finsler space equipped with fundamental metric function $F(x, \dot{x})$ satisfying the requisite condition for conditions for being a Finsler metric and $g_{ij}(x, \dot{x})$ be the fundamental metric tensor defined by $F^2(x, \dot{x}) = g_{ij}(x, \dot{x})\dot{x}^i\dot{x}^j$. Let $T_j^i(x, \dot{x})$ be any tensor field, the covariant derivative of such a tensor field in the sense of cartan is given by Rund [2]

$$(1.1) \quad T_{j|k}^i = \partial_k T_j^i - \dot{\partial}_k T_j^i \dot{\partial}_k G^h + T_j^m \Gamma_{mk}^{xi} - T_m^i \Gamma_{jk}^{xm}$$

where $\Gamma_{jk}^{xi}(x, \dot{x})$ are the cartan connection coefficients satisfying the relation

$$(1.2) \quad (a) \Gamma_{jk}^{xi} = \Gamma_{kj}^{xi}, \quad (b) \dot{\partial}_h \Gamma_{jk}^{xi} \dot{x}^h = 0$$

(1.3)

We are quoting the following commutation formulae which shall be used in the later discussion:

$$(1.4) \quad T_{j|hk}^i - T_{j|kh}^i = -\dot{\partial}_r T_j^i K_{mhk}^r \dot{x}^m + T_j^r K_{rhc}^i - T_r^i K_{jkh}^r,$$

$$(1.4) \quad (\dot{\partial}_k T_j^i)_{|h} - \dot{\partial}_k T_{j|h}^i = \dot{\partial}_r T_j^i C_{hklm}^r \dot{x}^m - T_j^r \dot{\partial}_k \Gamma_{rh}^{xi} + T_r^i \dot{\partial}_k \Gamma_{jh}^{xr}$$

Where curvature tensor field $K_{jkh}^i(x, \dot{x})$ is given by

$$(1.5) K_{jkh}^r = Q_{(h,k)} \left\{ \frac{\partial \Gamma_{jh}^{xi}}{\partial x^k} - \frac{\partial \Gamma_{jh}^{xi}}{\partial x^r} G_k^r + \Gamma_{mk}^{xi} \Gamma_{jh}^{xm} \right\}$$

Where $Q_{(h,k)}\{\dots\dots\dots\}$ stands for the interchange of the indices h and k and subtraction thereafter. The curvature tensor field $K_{jkh}^i(x, \dot{x})$ is a homogenous function of degree zero in its directional arguments. The term $C_{jk}^i(x, \dot{x})$ appearing in (1.4) is any tensor field give by the following equation:

$$(1.6) C_{ijk}(x, \dot{x}) = \frac{1}{2} \frac{\partial g_{ij}(x, \dot{x})}{\partial \dot{x}^k} = \frac{1}{4} \frac{\partial^3 F^2(x, \dot{x})}{\partial \dot{x}^i \partial \dot{x}^j \partial \dot{x}^k}$$

We now give the following definition which shall be used in the later discussions.

If the recurrent normal projective curvature tensor of a Finsler space be assumed to be decomposable in the form (1.6) then such a space is always normal projective recurrent.

We now consider the last case in which the normal projective curvature tensor of the Finsler space is decomposable in the form (1.5). Transvecting (1.6) by x^j and thereafter using (1.4), we get

$$(2.1) \quad H_{kh}^i = X_k^i x^j Y_{jh}$$

Transvecting (2.1) by y_i , we get

$$(2.2) \quad X_j^i x^j y_i Y_{kh} = 0,$$

Form (2.2) we conclude that either of the following two conditions will always hold the space under consideration

$$(2.3) (a) X_k^i x^j y_i = 0, \quad (b) Y_{kh} = 0$$

If the condition (2.3b) automatically leads to $N_{jkh}^i = 0$, therefore such a condition is always not possible. Hence, only alternative left with us is to consider the case:

$$(2.4) X_j^i x^j y_i = 0$$

Contracting (2.1) with respect to the indices i and j we get

$$(2.5) H_{rkh}^r = H_{hk} - H_{kh} = X_r^r Y_{kh}$$

From (2.5), we have

$$(2.6) Y_{kh} = \frac{1}{X} (H_{hk} - H_{kh}) \quad \text{where } X = X_r^r$$

Therefore, we can state:

If the normal projective curvature tensor of a Finsler space be assumed to be decomposable in the form (2.7) then the tensors X_j^i and Y_{kh} always satisfy (2.4) (2.5) and (2.6).

REFERENCES

1. Prananovitch, M. *Projective and conformal transformation in recurrent and Ricci-recurrent Riemannian space, Tensor (N.S.), 12, 219-226, (1962).*
2. Rund, H. *The differential geometry of Finsler spaces, Springer Verlag, Springer Verlag, Berlin (1959).*
3. Sinha, B.B. & Singh, S.P. *On the recurrent tensor fields of a Finsler space, Rev. dela Fac. de, Sc, de. Istambul, Ser. A. Vol 33, 114-117, (1968).*
4. Sinha, R.S. *Infinitesimal projective transformation in a Finsler space, Prog. Maths, 5(1-2), 30-34, (1971).*
5. Yano, K. *Theory of Lie-derivatives and its applications, North Holland Publ. Co. Amsterdam (1957).*